

A remark on the matrix Airy function

A.M. Perelomov *

*Max-Planck-Institut für Mathematik,
Vivatsgasse 7, 53111 Bonn, Germany***Abstract**

An integral representation for matrix Airy function is presented.

In memory of M.V. Saveliev

The well-known Witten conjecture [1], that the logarithm of the partition function of general one-matrix model can be considered as the generating function for intersection numbers on the moduli space of algebraic curves, was proved by Kontsevich [2,3], who reduced this problem to the computation of the matrix Airy function (see [4–6], where many aspects of this problem were discussed). The matrix Airy function is defined by an integral representation. We give another integral representation for this function.

Following [3], we define the matrix Airy function on the space $\mathcal{H}_N$ of Hermitian $N \times N$ matrices by the integral

$$A(X) = \int \exp \left\{ i \left(\frac{1}{3} \operatorname{tr}(Y^3) - \operatorname{tr}(XY) \right) \right\} dY \quad (1)$$

over the space $\mathcal{H}_N$, where dY is a $U(N)$ -invariant measure on this space, $X, Y \in \mathcal{H}_N$. We also note that the function $A(X)$ satisfies the equation

$$\Delta A(X) + (\operatorname{tr} X) A(X) = 0, \quad (2)$$

* On leave of absence from Institute for Theoretical and Experimental Physics, 117259 Moscow, Russia. Current e-mail address: perelomo@dftuz.unizar.es

where Δ is the Laplacian.

We recall that the classical Airy function

$$\text{Ai}(x) = \int_{-\infty}^{+\infty} \exp(i(y^3/3 - xy)) dy \quad (3)$$

is the unique solution (up to a scalar factor) of the differential equation

$$\text{Ai}''(x) + x \text{Ai}(x) = 0 \quad (4)$$

that is bounded on real axes. It follows from Eq.(1) that the dependence of $A(X)$ on the variables $\tilde{X}$ and $\xi = N^{-1} \text{tr} X$, where $X = \tilde{X} + \xi I$ and $\text{tr} \tilde{X} = 0$, can be separated. Indeed we have

$$X = \tilde{X} + \xi I, \quad Y = \tilde{Y} + \eta I, \quad \text{tr} \tilde{X} = \text{tr} \tilde{Y} = 0,$$

$$\text{tr}(Y^3) = \text{tr}(\tilde{Y}^3) + 3\eta \text{tr}(\tilde{Y}^2) + N\eta^3,$$

$$\text{tr}(XY) = \text{tr}(\tilde{X}\tilde{Y}) + N\xi\eta.$$

Theorem 1. *The integral representation*

$$\begin{aligned} A(X) &= A(\tilde{X}, \xi) = A(Q, \xi) \\ &= CN^{-1/3} \int \exp\left\{\frac{i}{3} \text{tr}(P^3)\right\} \text{Ai}\left(N^{-1/3}(N\xi - \text{tr}(P^2))\right) \Phi(Q|P) d\mu(P) \end{aligned} \quad (5)$$

is valid for matrix Airy function (1), where Q and P are diagonal matrices, $\tilde{X} = UQU^{-1}$, $\tilde{Y} = U_1PU_1^{-1}$, U and U_1 are unitary matrices, $\Phi(Q|P)$ is the zonal spherical function

$$\Phi(P|Q) = \frac{\det\{\exp(i(q_j p_k))\}}{V(P)V(Q)}. \quad (6)$$

$$Q = \text{diag}(q_1, \dots, q_N), \quad P = \text{diag}(p_1, \dots, p_N), \quad d\mu(P) = (V(P))^2 dP, \quad (7)$$

$C = \text{const}$, and $V(P)$ is the Vandermonde determinant.

$$V(P) = \prod_{j < k} (p_j - p_k) = \det(p_j^{k-1}). \quad (8)$$

Proof. This follows immediately from the Harish-Chandra formula.

Theorem 2 [7]. *If Φ is a conjugacy invariant function on the space Hermitian $N \times N$ matrices, then*

$$\begin{aligned} & \int \Phi(Y) \exp(-i \operatorname{tr}(QY)) dY \\ &= (-2\pi i)^{N(N-1)/2} (V(Q))^{-1} \int \Phi(P) \exp(-i \operatorname{tr}(QP)) (V(P))^2 dP \end{aligned} \quad (9)$$

for any diagonal Hermitian matrix Q , where $Y = UPU^{-1}$ and last integral is taken over the space of diagonal Hermitian matrices P .

We thus obtain an integral representation of the matrix Airy function such that the dependence on the variable ξ is contained only in the standard one-dimensional Airy function. This representation may be useful for investigation of properties of the matrix Airy function.

For the case $N = 2$, we have $\operatorname{tr}(P^3) = 0$, and integral representation (5) takes the simpler form

$$A(X) \equiv A(\xi, r) = \int \exp \left\{ \frac{2i}{3} (\eta^3 + 3\eta p^2 - 3\xi\eta) \right\} \Phi(r|p) p^2 dp d\eta, \quad (10)$$

where

$$\Phi(r|p) = \frac{\sin pr}{pr} \quad (11)$$

is the zonal spherical function for the matrix space with $N = 2$. The integration over η gives

$$A(\xi, r) = C \int \operatorname{Ai} \left(\left(\frac{3}{2} \right)^{1/3} (\xi - p^2) \right) \Phi(r|p) p^2 dp, \quad C = 2 \left(\frac{3}{2} \right)^{1/3}. \quad (12)$$

If we take the integral over p in (10), we obtain another integral representation

$$A(\xi, r) = C_1 \int \exp \left\{ \frac{2i}{3} (\tau^3 - 3\xi\tau) \right\} G(r, \tau) d\tau, \quad C_1 = \text{const}, \quad (13)$$

where

$$G(r, \tau) = \tau^{-3/2} \exp\{i\tau^{-1}r^2\} \quad (14)$$

is the Green function of the Schrödinger equation for free motion in three-dimensional space.

Acknowledgments. This note was written during a stay at Max-Planck-Institute für Mathematik, Bonn. It is pleasure to thank the staff of the institute for their hospitality.

References

- [1] Witten E., *Two-dimensional gravity and intersection theory on moduli space*, Surveys in Diff. Geometry (J. Differ. Geom. Suppl.) **1**, 243-310 (1991)
- [2] Kontsevich M., *Intersection theory on the moduli space of curves*, Func. Anal. Appl. **25**, No 2, 123 - 129 (1991)
- [3] Kontsevich M., *Intersection theory on the moduli space of curves and the matrix Airy function*, Commun. Math. Phys. **147**, 1-23 (1992)
- [4] Witten E., *On the Kontsevich model and other models of two-dimensional gravity*, Preprint IASSNS HEP-91/24, Princeton Univ., Princeton, NJ (1991)
- [5] Witten E., *The N matrix model and gauged WZW models*, Nucl. Phys. **B371**, 191-245 (1992)
- [6] Itzykson C. and Zuber J.-B., *Combinatorics of the modular group.II: The Kontsevich integrals*, Int. J. Mod. Phys. **A7**, 5661-5705 (1992)
- [7] Harish-Chandra, *Differential operators on semisimple Lie algebra*, Amer. J. Math. **79**, 87-120 (1957)